

SPIT OR SWALLOW

Cum in the mouth increases the risk of HIV being passed on. There seems to be little difference between spitting and swallowing. If you're worried about sucking or being sucked it's worth knowing that most men get HIV from unprotected fucking and not unprotected sucking. We think that most men who've been infected with HIV from sucking got cum in their mouths.

Cock-sucking: use your head

Terrence
HIGGINS
TRUST



SAY AAAAH

Throat infections increase the risk of getting HIV. If you're worried about sucking cock it's worth knowing that most men get HIV from unprotected fucking and not unprotected sucking. The risk increases if you get cum in your mouth, especially if you have a sore throat or infection. But some sexual infections can affect your throat and don't feel painful. You can get these checked out at a sexual health clinic.

Cock-sucking: use your head

Terrence
HIGGINS
TRUST



OPEN WIDE



A healthy mouth reduces the risk of getting HIV. If you're worried about sucking cock it's worth knowing that most men get HIV from unprotected fucking and not unprotected sucking. The risk increases if you get infected cum in your mouth, especially if your gums bleed, you brush your teeth before sex or you have a mouth ulcer.

Cock-sucking: use your head

Terrence
HIGGINS
TRUST



SHOOTING YOUR LOAD

A high viral load can increase the risk of passing on HIV. If you have HIV, your viral load can rise when you have another infection or when you are run down. If you're worried about being sucked it's worth knowing that most men get HIV from unprotected fucking and not unprotected sucking. The risk increases if you cum in his mouth.

Cock-sucking: use your head

Terrence
HIGGINS
TRUST

